

**TABLE 3**  
**NEW YORK OFFICIAL “WITHDRAWAL  
TIMES”**

<b>Drug Name (trade name)</b>	<b>RCI Class</b>	<b>“Withdrawal Time”</b>
<b>Furosemide</b> (250 mg-500 mg administered on the racing grounds by a licensed racing association member.)	n/a	4-4.5 hours prior to post
<b>Aminophylline</b>	3	48 hours
<b>Anabolic Steroids</b> (Equipoise)	4	48 hours
<b>Arsenic Solution</b> (Fowlers Solution)	n/a	48 hours
<b>Aspirin or Sodium Thiosalicylate</b>	4	48 hours
<b>Biologics</b> (bacterins, antitoxins except tetanus antitoxin) They may not be administered within 48 hours of the start of a racing program. In this regard, substances ingested by a horse shall be deemed administered at the time of eating and drinking. It shall be part of the trainer’s responsibility to prevent such ingestion within such 48 hours.	n/a	48 hours
<b>Chymotrypsin</b> (Kymar)	n/a	48 hours
<b>Diuretics</b> (read furosemide rule)	n/a	48 hours
<b>Epinephrine</b> (Adrenaline)	2	48 hours
<b>Glucocorticoids</b> (Prednisolone)	4	48 hours
<b>Griseofulvin</b> (Fulvicin)	n/a	48 hours
<b>Guaiacol derivatives</b> (Guifenesin, Ripercol-L)	4	48 hours
<b>Hormones &amp; Steroids</b> (testosterone, progesterone, estrogen, chorionic gonadatropin, glucocorticoids [Prednisolone, Depomedrol], and anabolic steroids [Equipoise]) except in conjunction with joint aspiration as restricted in subdivision (f)*.	4	48 hours
<b>Hydroxychloroquine Sulfate</b> (Rheaform)	n/a	48 hours
<b>Iodine Injection</b> (Hypodermin, Harvey’s Injectible Blister)	n/a	48 hours
<b>NSAID’s: Phenylbutazone</b> (Butazolidin), <b>Flunixin</b> (Banamine), <b>Meclofenamic Acid</b> (Arquel), <b>Naproxen</b> (Naprosyn, Equiproxen)	4	48 hours
<b>Methenamine</b> (Urotropin)	n/a	48 hours
<b>Orgotein</b> (Palosein)	n/a	48 hours
<b>Sarapin</b>	n/a	48 hours
<b>Selenium/ Vitamin E (E-Se)</b>	n/a	48 hours

<b>Sodium Thiosalicylate</b>	n/a	48 hours
<b>Sulfonamide Drugs</b>	n/a	48 hours
<b>Theophylline</b>	3	48 hours
<b>Thiazide derivatives (Diuril)</b>	n/a	48 hours
<b>Trichlormethiazide &amp; Dexamethazone (Naquazone bolets)</b>	4	48 hours
<b>Antihistamines</b>	3?	72 hours
<b>Ketamine HCl</b>	2	72 hours
<b>Methocarbamol (Robaxin)</b>	4	72 hours
<b>Pentazocine (Talwin)</b>	3	72 hours
<b>Vermifuges (except phenothiazine)</b>	n/a	72 hours
<b>Xylazine (Rompum)</b>	3	72 hours
All other medications including procaine	n/a	168 hours (7 days)
Joint aspiration with steroid injection	n/a	120 hours

\*(f) No other drugs or medications (including procaine) may be administered by any means within one week of the start of a racing program. In addition, a horse which has had a joint aspirated (in conjunction with a steroid injection) may not race for at least five days following such procedure, and whenever such procedure is performed, the trainer shall notify the stewards of such fact, in writing, before the horse is entered to race.